



Buttocks Augmentation

Allows reshaping of flat or poorly developed buttocks to make them more beautiful, rounded and firmly shaped.

Price:

Buttocks Implant:	4,000 US\$ (inclusive of 3 nights hospital stay)
Lipofilling:	1,000 US\$

Pre- Op Instructions:

- Consultation with surgeon
- Evaluation for other medical conditions
- No intake of food or water 8 hrs prior to surgery
- Compression garment should be available to be used post-op
- Stop smoking & alcohol intake at least 2 weeks before surgery
- Anticoagulants, vitamins, hormones (including birth control pills), anti-depressants & some other medications also to be stopped at least 2 weeks before surgery.

NOTE: Please provide Yanhee with a complete list of medications each patient is taking so proper advice can be given as to which medications should or shouldn't be stopped.

Anesthesia: General anesthesia

Duration: About 2 hours



- NOTE: 1. Prices are subject to change without prior notice.
2. Further details of procedures may be requested through e-mail.

Website: www.yanhee.net E-mail: inter_contact@yanhee.net, info@yanhee.net
Hotline: (+66) 02-879-0300 ext. 7 (available 7AM-10PM Bangkok Local Time)

Procedure:

Buttocks Lipofilling

- Needle aspiration removes the patient's own fat from areas such as the abdomen, or thighs.
- With the use of a microcannula this fat is then injected into the buttocks. No resultant scar is expected.

Buttock Implant

Round, cohesive silicone gel implants

Brand: Sebbin – Made in France

- A 5-6 cm incision is made overlying the tailbone.
- A packet is made between the gluteal muscles.
- The silicone implant is inserted into the packet.
- The incision is sutured.

Post-op Instructions:

- Pressure dressing and garments should be worn for at least 2-3 days.
- The patient should lie prone for the first 3 days following surgery to avoid too much pressure on the wound.
(NOTE: This procedure may not be done with surgeries such as breast surgery, Tummy Tuck, etc.)
- Sitting should be avoided during the first 3 days. May sit alternately on one buttock on the 4th day. Normal sitting is allowed 1 week after surgery.
- Buttocks massage is not necessary.
- May return to work as follows:
 - 1-2 weeks – for clerical/office jobs
 - 2 weeks – for jobs requiring excessive walking
- Full physical activities can be gradually resumed 1-3 months after surgery.

IMPORTANT:

- Buttock Lipofilling – results will only last for 3-6 months as transferred fat is reabsorbed by the body.
- Buttock Implants – injections to the buttocks should be avoided in the future.

BEFORE



AFTER

